



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

enlarging on this subject, till another time ; and conclude with assuring you of the most perfect esteem, with which I have the honour to be,

Dear Sir,

Your most obedient,
humble servant,

Joseph Gaertner, M. D.

XIV. *A Catalogue of the Fifty Plants from Chelsea Garden, presented to the Royal Society by the worshipful Company of Apothecaries, for the Year 1760, pursuant to the Direction of Sir Hans Sloane, Baronet, Med. Reg. & Soc. Reg. nuper Præses, by John Wilmer, M. D. clariss. Societatis Pharmaceut. Lond. Socius, Hort. Chelsean. Præfectus & Prælector. Botanic.*

Read Feb. 19, } 1901
1761.

- A** LCEA vulgaris major, flore
ex rubro roseo. C. B. 316.
1902 Andromeda pedunculis aggregatis, corollis
cylindricis, foliis alternis ovatis integerrimis.
Linn. Spec. Plant. 293.
1903 Aristolochia longa vera. C. B. 107. Offic. 47.
1904 Asclepias foliis revolutis linearibus verticillatis,
caule erecto. Linn. Sp. 217.
1905 Asphodelus foliis planis, caule ramoso, floribus
sparsis. Dict. Hort. Icon.

- 1906 *Bacteria foliis ovatis oppositis, floribus lateralibus, caule fruticoso ramoso.* Dict. Hort.
- 1907 *Camphorosma foliis hirsutis linearibus.* Amæn. Acad. Sp. 392.
Camphorata hirsuta. C. B. P. 486.
- 1908 *Cardiaca.* J. B. 3. 56.
Marrubium Cardiaca dictum. C. B. 230.
Offic. 104.
- 1909 *Cedrus folio cupressi, media, majoribus baccis.*
C. B. P. 487.
Cedrus Phœnicia, altera Plinii et Theophrasti.
Lobel. 221.
- 1910 *Cistus arborescens foliis ovato lanceolatis acuminatis trinerviis sessilibus utrinque villosis.*
- 1911 *Coronilla fruticosa stipulis subrotundis.* Linn.
Sp. Pl. 743.
- 1912 *Coronopus sylvestris hirsutior.* C. B. 190.
Offic. 147.
- 1913 *Damafonium stellatum.* Lugd. 1058.
Plantago aquatica stellata. C. B. 190.
- 1914 *Filix mas non ramosa dentata.* C. B. 358.
- 1915 *Galeopsis palustris Betonicæ folio.* Tourn.
Panax Coloni Officin. 341.
- 1916 *Gentiana palustris angustifolia.* C. B. 188.
Pneumonanthe. Ger. 355.
- 1917 *Gundelia foliis pinnatifidis spinosis, capite araneosa lanugine obsita.* Miller's Icons.
- 1918 *Helleborine latifolia montana.* C. B. P. 186.
- 1919 *Hibiscus foliis inferioribus trilobis, summis quinque partitis obtusis crenatis, calycibus inflatis.* Dict. Hort.
- 1920 *Kalmia foliis lanceolatis corymbis lateralibus.*
Linn. Gen. nov. 1079.

- 1921 *Lycium foliis cordato-ovatis seffilibus oppositis perennantibus, spinis crassiss bigeminis, floribus confertis.* Dict. Hort.
- 1922 *Lycopodium Sabinæ facie.* Fl. Jenens. 328.
Muscus clavatus foliis cupressi. C. B. 360.
- 1923 *Lygusticum quod Sefeli Offic.* C. B. 162.
Offic. 452.
Siler montanum major. Mor. Umb. 7. 8.
- 1924 *Lyfimachia speciosa quibusdam onagra dicta filiquosa.* J. B. 2. 906.
Chamenerion flore Delphinii. Park. Par. 270.
- 1925 *Malva caule erecto herbaceo, foliis lobatis, spicis secundis axillaribus.* Linn. Sp. 688.
- 1926 *Magnolia foliis ovato-lanceolatis subtus glaucis annuis.* Dict. Hort.
- 1927 *Mespilus Cretica folio circinato et quasi cordiformi.* T. Cor. 43.
Chamæcerasus Idæa, Alpini Exotic. 5.
- 1928 *Mespilus folio subrotundo, fructo rubro.*
Tourn. 642.
- 1929 *Milium semine luteo vel albo.* C. B. 26.
Offic. 317.
- 1930 *Morus foliis palmatis, fructibus hispidis.* Linn. Sp. Pl. 986.
Morus sativa foliis urticæ mortuæ, cortice papyrifera. Kempf. Amæn. 471.
- 1931 *Narcissus spatha uniflora nectario maximo limbo fimbriato petalo longiore.*
- 1932 *Narcissus spatha uniflora nectarii limbo campanulato erecto petalo æquali.* Linn. Sp. 289.
- 1933 *Nasturtium supinum capsulis verrucosis.* Ray Method. emendata, 98.
Coronopus Ruellii. Ger. 346. Offic. 343.

- 1934 *Ornithopodium radice nodosa*. Park. 1093.
- 1935 *Panicum Germanicum* five *panicula minore*.
C. B. 27. Offic. 343.
- 1936 *Pimpinella sanguisorba minor*. C. B. 160.
Offic. 366.
- 1937 *Polypodium vulgare*. C. B. 359.
Polypodium quercinum. Offic. 379.
- 1938 *Ranunculus ceratophyllus*, *feminibus falcatis*
in spicam adactis. Mor. Hist. 2. 440.
- 1939 *Rhus foliis ternatis lineari-lanceolatis*, *integer-*
rimis petiolatis utrinque glabris.
- 1940 *Salvia foliis cordatis obtusis crenatis subtomen-*
tofis, *corollis calyce angustioribus*. Linn.
Sp. Pl. 25.
- 1941 *Scordium alterum*, five *Salvia sylvestris*.
Scorodonia. Offic. 438. C. B. 247.
- 1942 *Scrophularia major*. Park. 610.
Scrophularia nodosa foetida. C. B. P. 235.
Offic. 440.
- 1943 *Scrophularia aquatica major*. C. B. P. 235.
Betonica aquatica. Ger. 579. Offic. 441.
- 1944 *Solanum caule inermi subfruticoso*, *foliis ob-*
longo ovatis sinuatis utrinque glabris floribus
alaribus. Miller's Icons.
- 1945 *Sisarum Germanorum*. C. B. 155. Off. 456.
- 1946 *Sium arvense* five *segetum*. Tourn. 308.
Selinum Si foliis. Ger. emac. 1018.
- 1947 *Tormentilla sylvestris*. C. B. 326. Off. 489.
- 1948 *Valeriana floribus tetrandis æqualibus*, *foliis*
pinnatifidis, *feminibus palea ovali adnatis*.
Hort. Upsal. 13.
- 1949 *Veratrum flore subviridi*. Tourn. 272. Offic.
226.

1950 *Viburnum foliis ovatis acuminatis ferratis venosis, petiolis lævibus.*

XV. *An Account of the Cicuta, recommended by Dr. Storke; by William Watſon, M. D. F. R. S.*

To the Royal Society.

Gentlemen,

Read Feb. 17, 1761. **I**N a paper I lately laid before you, I endeavoured to demonstrate, that the *Cicuta major*, which, ſince the publication of Dr. Storke's work at Vienna, had been uſed medicinally in England, was the plant intended by that gentleman; and not the *Cicuta aquatica*, as had been ſuggeſted by ſome practitioners here. And Dr. Storke has removed every doubt, which could remain, by transmitting hither to Mr. Hudſon, a very ingenious apothecary and botaniſt, ſome leaves of the *Cicuta major*, or common hemlock, which grew at Vienna, and is of the ſame ſpecies with the plant ſo denominated here.

As Dr. Storke informs us, that, ſince the publication of his treatiſe, he has received letters from almoſt every part of Europe, confirming his good opinion of the virtues of the *Cicuta*, and as he is about to publiſh a ſecond treatiſe upon the ſame ſubject, containing ſtill more extraordinary relations of cures brought about, by adminiſtring that plant; there is no doubt therefore, but that endeavours will be